

Note

On the Spectrum of n Quasigroups with Given Conjugate Invariant Subgroup

D. G. HOFFMAN

Department of Mathematics, Auburn University, Auburn, Alabama 36849

Communicated by the Managing Editors

Received April 20, 1982

We show in this paper that for any group G of permutations of an $(n + 1)$ set, any composite $q \geq (n + 1)^2$ is the order of an n quasigroup Q whose conjugate invariant subgroup is precisely G .

If n is a positive integer, and X is a finite set of cardinality $|X| = q$, we say that Q is an n quasigroup on X of order q if Q is a $(q^n) \times (n + 1)$ array, with entries from X , with the property that each of the $n + 1$ $q^n \times n$ subarrays of Q consists of q^n distinct rows. Thus each of the q^n possible n tuples with entries from X occurs exactly once as a row in such a subarray. We consider two n quasigroups on X *equivalent* if a permutation of the rows of one produces the other.

If Q is an n quasigroup on X , we denote by $\Gamma(Q)$ the set of permutations of the columns of Q which produce an array equivalent to Q . Evidently, $\Gamma(Q)$ is a subgroup of the symmetric group S of permutations of $\{0, 1, \dots, n\}$. We call $\Gamma(Q)$ the *conjugate invariant subgroup* of Q . These have been studied previously (see, for example, [6, 7]).

For each subgroup G of S , we define $\Lambda(G)$, the *spectrum* of G , to be the set of all positive integers q for which there exists an n quasigroup Q of order q with $\Gamma(Q) = G$.

We are concerned here with the following:

Conjecture. For each subgroup G of S , $\Lambda(G)$ consists of all but finitely many positive integers.

If $\Gamma(Q) = S$, the n quasigroup Q is said to be *totally symmetric*. These exist for all orders; for example, take for the rows of Q the $(n + 1)$ tuples summing to zero in an Abelian group X . Thus $\Lambda(S)$ contains all positive integers, so the conjecture is true if $G = S$.

Lindner and Steedly [2] have established the conjecture for $n = 2$, i.e.,

$q^2 \times 3$ orthogonal arrays. McLeish has established the conjecture for $n = 3$ in most cases in [3–5], as well as the case $|G| = 1$ for all n . We showed in [1] that $A(G)$ is infinite if G is a semiregular group of permutations.

Our main result is:

THEOREM. *Let m and p be positive integers with $m > n$ and $p \geq 2$. Then for any subgroup G of S , there is an n quasigroup Q of order mp with $\Gamma(Q) = G$. In particular, every composite $q \geq (n + 1)^2$ is in $A(G)$.*

Proof. Let M and P be finite Abelian groups of orders m and p , respectively. We construct an n quasigroup Q on $X = M \times P$ as follows: Choose $n + 1$ distinct elements a_0, a_1, \dots, a_n in M , choose a pair b, c of distinct elements of P , and let $s = a_0 + a_1 + \dots + a_n$.

Now let $x_0, x_1, \dots, x_n \in M$, $y_0, y_1, \dots, y_n \in P$. We declare the $(n + 1)$ tuple $((x_0, y_0), (x_1, y_1), \dots, (x_n, y_n))$ to be a row of Q if and only if the following two conditions hold:

- (1) $x_0 + x_1 + \dots + x_n = s$;
- (2) if for some $\pi \in G$, $(x_0, x_1, \dots, x_n) = (a_{\pi(0)}, a_{\pi(1)}, \dots, a_{\pi(n)})$, then $y_0 + y_1 + \dots + y_n = b$. Otherwise, $y_0 + y_1 + \dots + y_n = c$.

It is easy to check that Q is an n quasigroup on X , and that $\Gamma(Q) = G$.

Finally, if $q \geq (n + 1)^2$ is composite, then q has a prime factor $p \leq \sqrt{q}$, and then $m = q/p > n$.

REFERENCES

1. D. G. HOFFMAN, On conjugate invariant subgroups of n quasigroups, in "Proc. 9th S-E Conf. Combinatorics, Graph Theory, and Computing," pp. 373–381.
2. C. C. LINDNER AND D. STEEDLEY, On the number of conjugates of a quasigroup, *Algebra Universalis* **5** (1975), 191–196.
3. M. MCLEISH, "On the Number of Conjugates of Ternary Quasigroups," Ph.D. Thesis, McMaster University, Hamilton, Ontario, 1976.
4. M. MCLEISH, On the number of conjugates of n -ary quasigroups, *Canad. J. Math.* **31** (3) (1979), 637–654.
5. M. MCLEISH, On the existence of ternary quasigroups with 2 or 8 conjugacy classes, *J. Combin. Theory Ser. A* **29** (2) (1980), 199–211.
6. A. SADE, Quasigroups parastrophiques, expressions et identities, *Math. Nachr.* **20** (1959), 73–106.
7. S. K. STEIN, On the foundations of quasigroups, *Trans. Amer. Math. Soc.* **85** (1957), 228–256.